

**Baxter**  
**Recothrom**  
THROMBIN TOPICAL  
[RECOMBINANT]

# 20,000 Unit\* Reconstitution Instructions

\* Units used herein represent international units of potency determined using a reference standard that has been calibrated against the World Health Organization Second International Standard for Thrombin.



Remove the contents of the **Recothrom** thrombin topical (recombinant) Reconstitution Kit.



Remove the flip-off cap from the top of the **Recothrom** thrombin vial and the diluent vial. Attach a needle-free transfer device (one each) to the **Recothrom** thrombin and diluent vials and snap them into place by placing the vial on a flat surface and attaching the transfer device straight into the center of the vial stopper.



Open the sterile, empty 20 mL syringe package and attach the 20 mL syringe to the needle-free transfer device on the diluent vial. Injection of air into the diluent vial may facilitate withdrawal of the diluent.



Draw up 20 mL of diluent from the vial into the syringe.



Remove the diluent-filled syringe from the diluent vial and attach it to the transfer device on the **Recothrom** thrombin vial.

Transfer the 20 mL of diluent from the syringe into the **Recothrom** thrombin vial; the vacuum in the vial facilitates transfer.



Leave the syringe attached and gently swirl and invert the **Recothrom** thrombin vial until the powder is completely dissolved. Avoid excessive agitation.



Remove pre-printed “**DO NOT INJECT**” label from the outside of the syringe package and apply it to the syringe.



Withdraw **Recothrom** thrombin solution.

**RECOTHROM** thrombin is now ready to be transferred to a labeled receptacle in a sterile field.

**RECOTHROM THROMBIN IS FOR TOPICAL USE ONLY – DO NOT INJECT.**

No **RECOTHROM** thrombin kit components contain latex. Use aseptic technique when handling the vials and syringes.

PLEASE CONSULT THE FULL DRUG PRESCRIBING INFORMATION FOR FULL DETAILS.

**Baxter**

**Baxter**  
**Recothrom**  
THROMBIN TOPICAL  
(RECOMBINANT)

# 5,000 Unit\* Reconstitution Instructions

\* Units used herein represent international units of potency determined using a reference standard that has been calibrated against the World Health Organization Second International Standard for Thrombin.



Remove the contents of the **Recothrom** thrombin topical (recombinant) Reconstitution Kit.



Remove flip-off cap from the top of the **Recothrom** thrombin vial. Attach the needle-free transfer device. Snap it into place on the vial by placing the vial flat on a surface and attaching the transfer device straight into the center of the vial stopper.



Attach the prefilled diluent syringe to the needle-free transfer device.



Inject the 5 mL of diluent from the syringe into the product vial. Keep the syringe plunger depressed.

**NOTE: DO NOT** reuse the diluent syringe for transfer of the reconstituted product. Remove and discard the diluent syringe.



Gently swirl and invert the product vial until the powder is completely dissolved. Avoid excessive agitation.



Remove pre-printed “DO NOT INJECT” label from the outside of the syringe package. Apply the label to the provided empty transfer syringe.



Attach transfer syringe to **Recothrom** thrombin vial and withdraw solution.

**RECOTHROM** thrombin is now ready to be transferred to a labeled receptacle in a sterile field.

**RECOTHROM THROMBIN IS FOR TOPICAL USE ONLY – DO NOT INJECT.**

No **RECOTHROM** thrombin kit components contain latex. Use aseptic technique when handling the vials and syringes.

The information presented here is based on the **Recothrom** thrombin Prescribing Information.

BXU601996 Baxter and Recothrom are trademarks of Baxter International Inc. or its subsidiaries.