

Preventing Falls

Optimal Bed Height

Falls are a problem

Falls are the most commonly reported hospital adverse event¹ in the adult inpatient setting and a single fall with serious injury costs the hospital on average \$13,000.² Evidence-based protocols and equipment that help prevent falls improve patient safety and reduce cost.

Rethinking low-low beds

Historically, low-low beds have been indicated as a falls reduction measure. The thought is that the patient will not be able to exit a very low bed, creating a form of passive restraint, and will fall a shorter distance if rolling or sliding out of the bed.³

Evidence from a large, randomized controlled trial of 10,937 patients across 18 hospital wards does not support this thinking. The number of falls in the bedroom per 1,000 occupied bed days did not differ significantly between hospital units using conventional beds and those with low-low beds (Table 1). In fact, the rates of falls in the bedroom and falls with injury were slightly higher in the units with low-low beds.³

Table 1: Median outcome rates comparing conventional and low-low beds³

<i>Per 1,000 occupied bed days</i>	<i>Units with low-low beds</i>	<i>Units without low-low beds</i>
Falls in the bedroom	2.2	1.7
Falls with injury	1.7	1.2

While the use of low-low beds may be an appropriate strategy in some cases, they present hazards to falls risk patients, including the elderly. Biomechanical patient-based studies demonstrate that entry and egress from a low-low bed increases postural demands, putting patients at greater risk for falling.⁴⁻⁶ One author concludes that the recommendation to use the lowest bed frame possible is 'inaccurate and unsafe, and is not based on patient performance.'⁶ A safe solution is to provide fall risk patients with beds that are within the optimal height range.

Determining optimal bed height range

A hospital bed should adjust to a range of heights to meet the capabilities of individual patients as they move from the bed. Available range of motion determines a patient's ability to rise from the edge of a bed or chair. Adjusting the height of the bed helps patients overcome these limitations.⁷

The optimal bed height for patients with sit-to-stand difficulties, including the elderly, is 120% of the patient's lower leg length. A height less than 120% makes sit-to-stand transitions more difficult for the patient by requiring a greater range of joint motion and velocity.⁷

Norms for lower leg length, as measured from the floor to the mid-patella, range from about 16.4 inches to 21.7 inches.⁸ To give the greatest number of patients the optimal bed height for their lower leg length, an inpatient bed needs to provide a range of heights from 19.7 inches to accommodate almost all women and up to 26.0 inches to accommodate almost all men, as shown in Table 2. The range of 19.7 to 26.0 inches for optimal bed height can be measured as the distance from the floor to the buttocks of the patient when seated. The thickness of the mattress will need to be taken into consideration when determining the optimal bed height, adding it to the height of the bed frame.

Preventing falls with optimal bed height
To help improve sit-to-stand performance and reduce falls risk and related serious injuries, beds need to offer an optimal seated height no lower than 19.7 inches to accommodate almost all women and up to 26.0 inches to accommodate almost all men. Hill-Rom's Med-Surg beds with surfaces are within the optimal bed height range and when used in conjunction with a falls protocol can help reduce patient falls related to bed ingress/egress.

Preventing Falls — Optimal Bed Height

Table 2: Lower leg length and optimal bed height^{7,8}

Range of optimal bed heights for sit-to-stand transition

Low of 19.7" to accommodate most women



High of 26.0" to accommodate most men

Lower leg length norms for women and men

5th percentile female 95th percentile male

Lower leg length* (100%)	16.4"	21.7"
Optimal bed height (120%)	19.7"	26.0"

*from floor to mid-patella



A bed height of about 120% of lower leg length facilitates standing which may reduce the frequency of falls in the inpatient setting.⁶ Bed heights significantly lower than 120% make it more difficult for the patient by requiring a greater range of motion to stand.

Successful Fall Prevention

Research shows that ingress and egress from low-low beds requires increased joint motion which can compromise balance, thus increasing the risk for patients that are already at risk for falls.⁴⁻⁶ Multicomponent fall reduction strategies including the use of the correct equipment and programs have shown to reduce falls by as much as 30%.⁹ Providing patients with beds within the optimal bed height helps reduce the risk of falling, thereby helping enhance patient safety when moving from the bed.

References

1. Currie L. 2008. *Patient Safety and Quality: An Evidence-Based Handbook for Nurses*.
2. Wong CA, et al. *The Joint Commission Journal on Quality and Patient Safety*. 2011;37(2):81-87.
3. Haines TP, Bell RA, Varghese PN. 2010 Pragmatic, Custer Randomized Trial of a Policy to Introduce low-low beds to hospital wards for the prevention of falls and fall injuries. *J Am Geriatr Soc*. 58(3):435-41.
4. Merryweather AS, Morse JM, Doig AK, Godfrey NW, Gervair P, Bloswick DS. 2015. Effects of bed height on the biomechanics of hospital bed entry and egress. *Journal of Work* 52(3):707-13.
5. Christman M, Morse J, Wilson C, Godfrey N, Doig A, Bloswick D, Merryweather A. 2015. Analysis of the influence of hospital bed height on kinematic parameters associated with patient falls during egress. *Procedia Manufacturing* 280-287.
6. Morse JM, Gervais P, Pooler C, Merryweather A, Doig AK, Bloswick D. 2015. The safety of hospital beds: ingress, egress and in-bed mobility. *Global Qualitative Nursing Research* 2:1-20.
7. Janssen WG, Bussmann HB, Stam HJ. 2002. Determinants of the sit-to-stand movement: A review. *Physical Therapy*. 82(9):866-879.
8. Gordon CC, Churchill T, Clouser CE, Bradtmiller B, McConville JT, Tebbets I, Walker RA. 1988 Anthropometric Survey of US Army Personnel: Methods and Summary Statistics. Technical Report TR-89-044 (AD A225 094). US Army Natick Research, Development, and Engineering Center, Natick, MA. Page 229, accessed online February 4, 2015: <http://www.google.com/url?sa=&rct=j&q=&esrc=s&source=web&cd=1&ved=0CCAQFjAA&url=http%3A%2F%2Fwww.dtic.mil%2Fcginbin%2FGetTRDoc%3FAD=ADA225094%26Location=U2%26doc=GetTRDoc.pdf&ei=dFTSVD7E4O5yQTazYHoDg&usg=AFQjCNHwCV552PqPxVv5xu1eT1PAFPzSQw&sig2=rGc>
9. Miale-Lye IM, Hempel S, Ganz DA, Shekelle PG. 2013. Inpatient fall prevention programs as a patient safety strategy: a systematic review. *Ann Intern Med* 158 (5 Pt 2):390-6.

Hill-Rom reserves the right to make changes without notice in design, specifications and models. The only warranty Hill-Rom makes is the express written warranty extended on the sale or rental of its products.

©2016 Hill-Rom Services, Inc. ALL RIGHTS RESERVED.
191351 rev 3 15-AUG-2016 ENG - US

For further information about this product or a service, please contact your local Hill-Rom representative or visit our webpage:

USA 800-445-3730
Canada 800-267-2337

www.hill-rom.com

Enhancing outcomes for
patients and their caregivers.

Hill-Rom